

Feshbach resonances in a strictly two-dimensional atomic Bose gas

K.K. Rajagopal, P. Vignolo^{*} and M.P. Tosi

*Classe di Scienze, Scuola Normale Superiore,
Piazza dei Cavalieri 7, I-56126 Pisa, Italy*

Abstract

We discuss the atom-atom scattering problem across a Feshbach resonance in a two-dimensional dilute Bose gas at zero temperature, in the limit where the s -wave scattering length exceeds the width of the vertical confinement. We determine a tunable coupling-strength parameter and by controlling it we evaluate how the condensate wave function spreads out with increasing atom-atom repulsions. We also discuss the stability of the condensate in the magnetic-field regime where the coupling has become attractive.

Key words: Bose gases, low dimensional, density profiles

PACS: 03.75.Hh, 32.80.Pj

1 Introduction

An ultracold two-dimensional (2D) Bose gas exhibits in several respects a different physical behaviour from its three-dimensional (3D) counterpart [1,2]. In the macroscopic limit phase fluctuations at any finite temperature destroy the long-range order associated with Bose-Einstein condensation, but still allow a Kosterlitz-Thouless transition to a superfluid “quasi-condensate” state characterized by local phase coherence [3]. A true condensate is restored in a trapped 2D gas by modifications induced by the confinement in the density of states [4], but the magnitude and sign of the scattering amplitude for atom-atom collisions can depend on external system parameters [5].

In fact, when the scattering length becomes comparable with the trap size a new two-particle scattering problem needs to be solved. The T-matrix for

^{*} Corresponding author, e-mail: vignolo@sns.it

two-body collisions in vacuo at low momenta and energy, which is used in 3D to obtain the collisional coupling parameter, vanishes in the strictly 2D limit as the axial width of the trapping potential becomes smaller than the s -wave scattering length [1,2]. It is then necessary to evaluate the scattering processes between pairs of Bose particles by taking into account the presence of the surrounding bosons through a many-body T-matrix formalism [6–8].

The 3D s -wave scattering length can be tuned by means of external electric or magnetic fields [9–11]. Within this process a resonance can appear at a particular value of the magnetic field and allow control of the coupling strength ranging from positive (repulsive) to negative (attractive) values. One may study new dynamical effects of collective oscillations in large condensates [12] and the dynamics of collapsing and exploding condensates [13]. Calculations of such Feshbach resonances have been given for 3D gases of ^7Li [14], ^{23}Na [10], ^{39}K and ^{41}K [15], ^{85}Rb and ^{87}Rb [11], and ^{133}Cs [9]. Experimental observations have been reported for ^{87}Rb [16], ^{85}Rb [13,17–19], ^{23}Na [20], and ^{133}Cs [21]. Feshbach resonances in quasi-2D atomic gases have been discussed by Wouters *et al.* [22], who have predicted a shift in the position of the resonance from the squeezing of the confinement.

In this work we examine the tuning of two-body collisions by means of a magnetic field in a strictly 2D condensate inside a harmonic trap at zero temperature. By modifying the coupling strength we can control the dimensionless physical parameter η which is the ratio between the mean-field interaction energy and the harmonic-oscillator level spacing [23]. In the case of repulsive interactions ($\eta > 0$), starting from a weakly interacting regime ($\eta \ll 1$) and up to a strongly interacting one ($\eta \gg 1$), we evaluate the equilibrium density profile as a function of η by solving numerically a non-linear Schrödinger equation by a split-step Crank-Nicholson discretization scheme. In the case of attractive interactions ($\eta < 0$) we can give an upper bound for the number of bosons which can condense without the condensate collapsing. At variance from the 3D gas (see for instance [24,25]), we have found that in 2D this critical number does not depend explicitly on the strength of the harmonic confinement.

The paper is organized as follows. Section 2 summarizes the Feshbach resonance formalism and applies it to the 2D Bose gas. In Sec. 3 we evaluate the equilibrium density profiles for the case of repulsive interactions and discuss the conditions for a stable condensate in the attractive coupling regime. Some concluding remarks are presented in Sec. 4.

2 Essential formalism

2.1 Overview on scattering theory

We start by recalling some well-known facts about a collision between two particles with internal degrees of freedom governed by an internal Hamiltonian H^{int} ,

$$H^{int} |\alpha\rangle = \epsilon_\alpha |\alpha\rangle \quad (1)$$

where $|\alpha\rangle$ and ϵ_α are the single-atom eigenstate and its corresponding eigenvalue. For example the internal Hamiltonian can describe the hyperfine interaction between nuclear and electronic spins and their Zeeman coupling to an external magnetic field B . A two-body collision is described by the Hamiltonian

$$H = \frac{p^2}{2m_r} + \sum_{i=1}^2 H_i^{int} + V(\mathbf{r}) \quad (2)$$

in the center-of-mass system, with $\mathbf{p}$ the relative momentum and m_r the reduced mass (see for instance [26]). The Hamiltonian in Eq. (2) is the sum of a part with eigenstates $|\alpha\beta\rangle$ (“channels”) tending asymptotically to a symmetrized product of separate-atom internal states $|\alpha\rangle$ and $|\beta\rangle$, and of a finite-range interaction $V(\mathbf{r})$ which couples the channels. The energy associated with an eigenstate $|\alpha\beta\rangle$ is $E_{\alpha\beta} = \epsilon_\alpha + \epsilon_\beta + \hbar^2 k_{\alpha\beta}^2 / 2m_r$, where $\hbar \mathbf{k}_{\alpha\beta}$ is the relative momentum of the two incoming particles. The asymptotic scattering state $\Psi_{\alpha\beta}(\mathbf{r})$ can be expanded on all scattering channels in the form

$$\Psi_{\alpha\beta}(\mathbf{r}) = \exp(i\mathbf{k}_{\alpha\beta} \cdot \mathbf{r}) |\alpha\beta\rangle + \sum_{\alpha',\beta'} f_{\alpha\beta}^{\alpha'\beta'}(\mathbf{k}_{\alpha\beta}, \mathbf{k}'_{\alpha'\beta'}) \frac{\exp(i\mathbf{k}'_{\alpha'\beta'} \cdot \mathbf{r})}{r} |\alpha'\beta'\rangle, \quad (3)$$

where the outgoing channel is denoted by $|\alpha'\beta'\rangle$ and the scattering amplitude $f_{\alpha\beta}^{\alpha'\beta'}$ is proportional to the T-matrix element $\langle \mathbf{k}'_{\alpha'\beta'} | T | \mathbf{k}_{\alpha\beta} \rangle$. The asymptotic magnitude of the momentum for the $|\alpha'\beta'\rangle$ channel is $k_{\alpha'\beta'} = \sqrt{2m_r(E_{\alpha\beta} - \epsilon_{\alpha'} - \epsilon_{\beta'})}$, from conservation of total momentum and energy in the scattering process. The channel is said to be open (closed) if the momentum takes positive real (imaginary) value.

The whole Hilbert space describing the spatial and spin degrees of freedom is now divided into a subspace P for open channels and a subspace Q including all closed channels [27]. The state vector $|\Psi\rangle$ and the Schrödinger equation $H|\Psi\rangle = E|\Psi\rangle$ are projected onto the two subspaces by means of projection operators P and Q satisfying the conditions $P + Q = 1$ and $PQ = 0$. The formal solution of the coupled equations obeyed by the two projected components is

$$|\Psi_Q\rangle = (E - H_{QQ} + i\delta)^{-1} H_{QP} |\Psi_P\rangle \quad (4)$$

and

$$(E - H_{PP} - H'_{PP}) |\Psi_P\rangle = 0, \quad (5)$$

where $H_{PP} = PHP$ etcetera,

$$H'_{PP} = H_{PQ}(E - H_{QQ} + i\delta)^{-1}H_{QP}, \quad (6)$$

and δ is a positive infinitesimal. The term H'_{PP} in Eq. (5) describes the Feshbach resonances. It represents an effective non-local, retarded interaction in the P subspace due to transitions to the Q subspace and back. The Hamiltonian $H_{PP} + H'_{PP}$ gives the effective atom-atom interaction in the open-channel subspace, the effects due to the existence of bound states being taken into account through the term H'_{PP} .

2.2 Feshbach resonances in the 2D coupling

A Feshbach resonance results when true bound states belonging to the closed-channel subspace match the energy of open channels: transient transitions may then be possible during the collision process [10]. Setting $H_{PP} = H_0 + U_1$ where H_0 includes the kinetic energy of relative motion and the internal Hamiltonian, the scattering matrix elements in the P subspace between plane-wave states with relative momenta $\mathbf{k}$ and $\mathbf{k}'$ read

$$\langle \mathbf{k}' | T | \mathbf{k} \rangle \simeq \langle \mathbf{k}' | T_1 | \mathbf{k} \rangle + \langle \Omega_{1,+} \mathbf{k}' | H'_{PP} | \Omega_{1,+} \mathbf{k} \rangle. \quad (7)$$

Here $T_1 = U_1 + U_1 G_0 T_1$ represents the scattering process if Q subspace is neglected and $\Omega_{1,+} = (1 - G_0 U_1)^{-1}$ is the wave operator due to the interaction potential U_1 , with $G_0 = (E - H_0)^{-1}$ [28]. In Eq. (7) the effect of closed channels has been included at first order in H'_{PP} . In the limit of zero relative velocity Eq. (7) leads to an expression for the coupling strength,

$$g = \tilde{g} + \sum_n \frac{|\langle \psi_n | H_{QP} | \psi_0 \rangle|^2}{E_{th} - E_n} \quad (8)$$

where the sum is over all the states $|\psi_n\rangle$ in Q subspace and we have set $|\Omega_{1,+} \mathbf{k}\rangle \simeq |\Omega_{1,+} \mathbf{k}'\rangle \equiv |\psi_0\rangle$. Here, $\tilde{g}$ is the coupling strength estimated in the absence of closed channels and E_{th} is the threshold energy for the state $|\psi_0\rangle$ at vanishing kinetic energy. Finally, when the threshold energy is close to the energy E_{res} of one specific bound state $|\psi_{res}\rangle$, this contribution will be dominant and the contributions from all other states may be included in $\tilde{g}$ through an effective non-resonant scattering length a_{nr} . For a strictly 2D condensate we get

$$g = \frac{4\pi\hbar^2/m}{\ln|4\hbar/\mu m a_{nr}^2|} + \frac{|\langle \psi_{res} | H_{QP} | \psi_0 \rangle|^2}{E_{th} - E_{res}}, \quad (9)$$

where μ is the chemical potential of the condensate.

Equation (9) is our main result. The non-resonant first term has been calculated from the two-body T-matrix for a binary collision occurring at low momenta and energy in the presence of a condensate, which fixes the energy of each colliding atom at the chemical potential [29,30]. The resonant second term can be tuned by exploiting the dependence of the energy denominator on external parameters and in particular on the strength of the applied magnetic field. Explicitly, if the energy denominator vanishes at a value B_0 of the field, then by expansion around this value we have

$$E_{th} - E_{res} \approx (\mathbf{m}_{res} - \mathbf{m}_\alpha - \mathbf{m}_\beta)(B - B_0) \quad (10)$$

where $\mathbf{m}_{\alpha,\beta} = -\partial\epsilon_{\alpha,\beta}/\partial B$ are the magnetic moments of the two colliding atoms in the open channel and $\mathbf{m}_{res} = -\partial E_{res}/\partial B$ is the magnetic moment of the molecular bound state. Equation (9) then yields

$$g = \frac{4\pi\hbar^2}{m} \left(\frac{1}{\ln |4\hbar/\mu m a_{nr}^2|} + \frac{\Delta B}{B - B_0} \right), \quad (11)$$

where the width parameter ΔB is defined as

$$\Delta B = \frac{m}{4\pi\hbar^2} \frac{|\langle \psi_{res} | H_{QP} | \psi_0 \rangle|^2}{\mathbf{m}_{res} - \mathbf{m}_\alpha - \mathbf{m}_\beta}. \quad (12)$$

Higher-order terms in H'_{PP} will lead to a broadening of the resonance. However, the width of resonant states close to the threshold energy in the open channel is usually small because of the low density of states, and Feshbach resonances are usually very sharp [28].

In the next Section we will analyze the consequences of tuning the coupling strength on the equilibrium state of a 2D condensate in a harmonic trap.

3 Equilibrium properties as functions of the coupling

We consider a Bose gas with a fixed number N of particles in a 2D harmonic potential $V_{ext}(r) = m\omega^2 r^2/2$. At zero temperature the condensate wave function $\Psi(r)$ is described by the mean-field Gross-Pitaevskii equation

$$\left[-\frac{\hbar^2}{2m} \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) + V_{ext}(r) + gn(r) \right] \Psi(r) = \mu \Psi(r), \quad (13)$$

where $n(r) = |\Psi(r)|^2$ is the density of the condensate. We introduce the dimensionless coordinate $x = r/a_{ho}$, where $a_{ho} = (\hbar/m\omega)^{1/2}$ is the harmonic oscillator length, and the dimensionless chemical potential $\mu' = 2\mu/\hbar\omega$. Thus

Eq. (13) can be written in rescaled form as

$$\left[-\frac{d^2}{dx^2} - \frac{1}{x} \frac{d}{dx} \right] \Psi(x) + x^2 \Psi(x) + 2\eta \left| \frac{\Psi(x)}{\Psi(0)} \right|^2 \Psi(x) = \mu' \Psi(x), \quad (14)$$

where $\eta = g |\Psi(0)|^2 / \hbar \omega$ is a dimensionless coupling parameter given by the ratio between the mean-field interaction energy per particle and the oscillator level spacing. The normalization condition on the wave-function is

$$N = a_{ho}^2 \int_0^\infty x dx |\Psi(x)|^2. \quad (15)$$

Equations (14) and (15) need to be solved self-consistently.

3.1 Repulsive coupling ($\eta > 0$)

In the case of repulsive interactions Eq. (14) always admits a solution: the condensate exists and is stable in the 2D harmonic trap. Of course, far away from the resonance the dimensionless coupling parameter η is primarily determined by the non-resonant scattering length. On approaching the resonance by increasing B towards B_0 , the coupling parameter becomes very small for $B \sim B_0 - \Delta B / (\ln |4\hbar / \mu m a_{nr}^2|)$: the non-linear term in Eq. (14) becomes small and the condensate wave function is close to a Gaussian profile. On further increasing B the gas enters the strong-coupling regime and in the limit $\eta \gg 1$ the kinetic energy term in Eq. (14) and the discrete structure of the trap levels become irrelevant. The profile then approaches a Thomas-Fermi form [31,32]. The shape of the condensate density profile, obtained by the numerical solution of Eq. (14) at several values of the coupling parameter η , is depicted in Fig. 1. The Figure illustrates comparisons with the free-gas and Thomas-Fermi profiles, according to the discussion given just above.

3.2 Attractive coupling ($\eta < 0$)

On going through the resonance the Thomas-Fermi-like density profile will suddenly shrink and collapse. Indeed, in the strong-coupling attractive regime the kinetic energy does not suffice to compensate for the negative mean-field energy term and Eq. (14) does not admit physical solutions. A condensate with a limited number of particles can only exist in the weakly attractive regime. An upper bound for the critical number N_c of bosons in a condensate with weakly attractive interactions can be estimated by comparing the kinetic energy and the mean-field energy at the centre of the trap. This leads to the inequality

$$1 + 2\eta \geq 0 \quad (16)$$




Fig. 1. Scaled density profile $|\Psi(x)|^2/|\Psi(0)|^2$ as a function of the dimensionless radial distance $x = r/a_{ho}$ for various values of the coupling parameter η . Dashed curves are Thomas-Fermi profiles and the dotted curve is the free-gas profile.

for condensate stability, where in the calculation of the kinetic energy we have used the free-gas wave function. Using the expression of η we find from Eq. (16)

$$N_c \simeq \left| \frac{1}{\ln |4\hbar/\mu m a_{nr}^2|} + \frac{\Delta B}{B - B_0} \right|^{-1}. \quad (17)$$

At variance from the trapped 3D gas [24], the critical number N_c does not depend on the harmonic oscillator length, the dependence being cancelled by the fact that in 2D the mean-field energy scales as $\approx a_{ho}^{-2}$ like the kinetic energy. Let us remark that the condensate in the case of attractive coupling can be metastable even if the number of condensed bosons is lower than N_c . A three-body loss term should be included in Eq. (13) in order to calculate the decay time and to describe the time evolution of such a condensate, as was done for the 3D Bose gas [25,33,34] and very recently for a 3D boson-fermion mixture [35]. An evaluation of three-body losses in the 2D Bose gas and of the time evolution of a 2D condensate in the weakly attractive coupling regime may be considered in the future.

4 Concluding remarks

In summary, we have applied the formalism already developed to treat Feshbach resonances in 3D boson gases to the case of a 2D Bose gas at zero temperature. One of the main results is that an external magnetic field could be used to drive the coupling from repulsive to attractive even in a pancake

geometry as the scattering length becomes larger than the axial width of the condensed cloud. For the case of repulsive interactions we have evaluated the particle density profiles at varying coupling strength, demonstrating the approach to a Thomas-Fermi profile followed by collapse. In the opposite regime of attractive coupling we have given an estimate for the critical number of bosons that can form a stable condensate. We have found that this critical number is inversely proportional to the absolute magnitude of the coupling, independently of the strength of the in-plane confinement.

Acknowledgements

The code used in the numerical calculations reported in Sec. 3.1 was originally developed by Professor B. Tanatar and his coworkers. We also acknowledge support from Scuola Normale Superiore di Pisa through an Advanced Research Initiative. One of us (MPT) thanks Professor V. E. Kravtsov and the Condensed Matter Theory group of the Abdus Salam ICTP for their hospitality during the final stages of this work.

References

- [1] M. Schick, Phys. Rev. A 3 (1971) 1067.
- [2] V. N. Popov, Functional Integrals in Quantum Field Theory and Statistical Physics, Chap. 6, Reidl, Dordrecht, 1983.
- [3] D. S. Fisher, P. C. Hohenberg, Phys. Rev. B 37 (1988) 4936.
- [4] V. Bagnato, D. Kleppner, Phys. Rev. A 44 (1991) 7439.
- [5] D. S. Petrov, M. Holzmann, G. V. Shlyapnikov, Phys. Rev. Lett. 84 (2000) 2551.
- [6] H. T. C. Stoof, M. Bijlsma, Phys. Rev. E 47 (1993) 939.
- [7] M. Bijlsma, H. T. C. Stoof, Phys. Rev. A 55 (1997) 498.
- [8] M. D. Lee, S. A. Morgan, M. J. Davis, K. Burnett, Phys. Rev. A 65 (2002) 043617.
- [9] E. Tiesinga, B. J. Verhaar, H. T. C. Stoof, Phys. Rev. A 47 (1993) 4114.
- [10] A. J. Moerdijk, B. J. Verhaar, A. Axelsson, Phys. Rev. A 51 (1995) 4852.
- [11] J. M. Vogels, C. C. Tsai, R. S. Freeland, S. J. M. F. Kokkelmans, Phys. Rev. A 56 (1997) 1067.

- [12] Y. Kagan, E. L. Surkov, G. V. Shlyapnikov, Phys. Rev. Lett. 79 (1997) 2604.
- [13] E. A. Donley, N. R. Claussen, S. L. Cornish, J. L. Roberts, E. A. Cornell, C. E. Weimann, Nature 412 (2001) 295.
- [14] P. O. Fedichev, Y. Kagan, G. V. Shlyapnikov, J. T. M. Walraven, Phys. Rev. Lett. 77 (1996) 2913.
- [15] H. M. J. M. Boesten, J. M. Vogels, J. G. C. Tempelaars, B. J. Verhaar, Phys. Rev. A 54 (1996) 3726.
- [16] N. R. Newbury, C. J. Myatt, C. E. Wiemann, Phys. Rev. A 51 (1995) 2680.
- [17] J. L. Roberts, N. R. Claussen, J. P. Burke, J. C. H. Green, E. A. Cornell, C. E. Weimann, Phys. Rev. Lett. 81 (1998) 5109.
- [18] P. Courteille, R. S. Freeland, D. J. Heinzen, F. A. van Abeelan, B. J. Verhaar, Phys. Rev. Lett. 81 (1998) 69.
- [19] S. L. Cornish, N. R. Claussen, J. L. Roberts, , E. A. Cornell, C. E. Weimann, Phys. Rev. Lett. 85 (2000) 1795.
- [20] S. Inouye, M. R. Andrews, H. J. Meisner, D. M. Stamper-Kurn, J. Stenger, H. J. Miesner, W. Ketterlee, Nature 392 (1998) 151.
- [21] V. Vuletic, A. J. Kerman, C. Chin, S. Chu, Phys. Rev. Lett. 82 (1999) 1406.
- [22] M. Wouters, J. Tempere, J. T. Devreese, cond-mat/0305235.
- [23] Y. Kagan, G. V. Shlyapnikov, J. T. M. Walraven, Phys. Rev. Lett. 76 (1996) 2670.
- [24] A. Eleftheriou, K. Huang, Phys. Rev. A. 61 (2000) 043601.
- [25] C. M. Savage, N. P. Robins, J. J. Hope, Phys. Rev. A 67 (2003) 014304.
- [26] H. T. C. Stoof, J. M. Koelman, B. J. Verhaar, Phys. Rev. B 38 (1988) 4688.
- [27] H. Feshbach, Ann. Phys. (NY) 19 (1962) 287.
- [28] C. J. Pethick, H. Smith, Bose-Einstein Condensation in Dilute Gases, Cambridge University Press, London, 2001.
- [29] S. A. Morgan, M. D. Lee, K. Burnett, Phys. Rev. A 65 (2002) 022706.
- [30] K. K. Rajagopal, P. Vignolo, M. P. Tosi, Physica B 344 (2004) 157.
- [31] V. V. Goldman, I. Silvera, A. Legget, Phys. Rev. B 24 (1981) 2870.
- [32] D. A. Huse, E. D. Siggia, J. Low Temp. Phys. 46 (1982) 137.
- [33] H. Saito, M. Ueda, Phys. Rev. A 65 (2002) 033624.
- [34] S. K. Adhikari, Phys. Rev. A 66 (2002) 013611.
- [35] S. K. Adhikari, cond-mat/0407032.